

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Figure 1 consists of 12 histograms arranged horizontally, labeled x_0 through x_{11} . Each histogram shows the frequency of non-zero elements in the vector x_k . The x-axis for each histogram is labeled x_k and ranges from 0 to 10. The y-axis is labeled 'Frequency' and ranges from 0 to 10. The distributions are roughly bell-shaped and centered around 5, with the peak frequency increasing from 10 at $k=0$ to 12 at $k=11$.

- 35



51

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R²⁴⁹ is Trp, (L) or (D)Lys, (L) or (D) Tyr or (D)Phe;

5 R²⁵⁰ is Arg;

R²⁵¹ is (L) or (D)Leu or Lys;

R²⁵² is (L) or (D)Arg;

R²⁵³ is (D)- or (L)- Phe;

R²⁵⁴ is Ala;

10 R²⁵⁵ is (D)- or (L)- Leu or is Lys;

R²⁵⁶ is absent or is (L) or (D) Arg;

R²⁵⁷ is (L) or (D) Tyr;

R²⁵⁸ is Ala; and

Y² is amide, thioether, thioester or disulfide.

15

8. The backbone cyclized analog of claim 7 wherein

R²⁴⁹ is Trp, (L)- or (D)- Lys or (D)Phe;

R²⁵⁰ is Arg;

R²⁵¹ is Lys or (D)Leu;

20 R²⁵² is (D)Arg;

R²⁵³ is (D)- or (L)- Phe;

R²⁵⁴ is Ala;

R²⁵⁵ is (D)- or (L)- Leu;

R²⁵⁶ is absent or is Arg;

25 R²⁵⁷ is (D)Tyr;

R²⁵⁸ is Ala; and

Y² is amide, thioether, thioester or disulfide.

30

9. The backbone cyclized IL-6 antagonist of claim 8 having the formula:


Trp-Arg-Lys-(D)Arg-Phe-AlaC3-Leu-Arg-(D)Tyr-AlaN3-NH₂

10. The backbone cyclized IL-6 antagonist of claim 8 having the formula:


35

(D)Lys-Arg-(D)Leu-(D)Arg-(D)Phe-AlaC3-(D)Leu-Arg-(D)Tyr-AlaN3- NH₂

11. The backbone cyclized IL-6 antagonist of claim 8 having the formula:



12. The backbone cyclized analog of claim 1 having the general formula 3:



Formula No. 3

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R¹ is (D)Bip, Gln, Lys, Lys(ZCL) or Dab;

R² is (D)Lys, Gly, Ala or Trp

R³ is Orn, 4PyrAla, (L) or (D)Dab, (D)Arg, Lys or Dpr;

R⁴ is Lys, Lys(ZCL), Arg, Arg(Mtr) or (D)Glu;


R⁵ is Asn, Trp or (D)Ala;

R⁶ is Arg, (p-NO₂)Phe, (L)- or (D)- Trp, Gln, Abu or Glu;

and

Y² is amide, thioether, thioester or disulfide.

13. The backbone cyclized analog of claim 1 having the general formula 4:



Formula No. 4

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R¹ is (D)Phe or Lys;


R² is (D)Cit, Lys or (D)Bip;

R³ is Dpr, 4PyrAla or (L)- or (D)- Arg;

R⁴ is HomArg, Orn or Lys;

R⁵ is (D)Gln or (L)- or (D)- Trp;
R⁶ is (L)- or (D)- Gln or (p-NO₂)Phe; and
Y² is amide, thioether, thioester or disulfide.

- 5 14. A pharmaceutical composition comprising a backbone
cyclized IL-6 antagonist comprising a peptide sequence of
five to twenty amino acids that incorporates at least one
building unit, said building unit containing one nitrogen
atom of the peptide backbone connected to a bridging group
10 comprising an amide, thioether, thioester or disulfide,
wherein the at least one building unit is connected via
the bridging group to form a cyclic structure, together
with a pharmaceutically acceptable carrier or diluent.
- 15 15. The pharmaceutical composition of claim 14 wherein the
IL-6 antagonist is a backbone cyclized peptide analog
having the general formula 1:



Formula No. 1

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R²⁴⁹ is Trp, / (L) or (D) Lys, (L) or (D) Tyr or (D) Phe;

R²⁵⁰ is Arg; /

R²⁵¹ is (L) or (D) Leu or Lys;

R^{252} is (L) or (D) Arg;

R^{253} is (D) - or (L) - Phe;

R²⁵⁴ is Ala;

R²⁵⁵ is/ (D)- or (L)- Leu or is Lys;

R²⁵⁶ is absent or is (L) or (D) Arg;

R²⁵⁷ is (L) or (D) Tyr;

R²⁵⁸ / is Ala; and

Y² is amide, thioether, thioester or disulfide.

16. The pharmaceutical composition of claim 15 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:

17. The pharmaceutical composition of claim 15 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:

18. The pharmaceutical composition of claim 15 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:

19. The pharmaceutical composition of claim 14 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the general formula 3:

Formula No. 3

X designates a terminal carboxy acid, amide or alcohol group;


R² is (D)¹Lys, Gly, Ala or Trp

R^4 is Lys, Lys(ZCL), Arg, Arg(Mtr) or (D)Glu;

R⁶ is Arg, (p-NO₂) Phe, (L) - or (D) - Trp, Gln, Abu or Glu;

Y^2 /is amide, thioether, thioester or disulfide.

having the general formula 4:



Formula No. 4

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R¹ is (D)Phe or Lys;

R² is (D)Cit, Lys or (D)Bip;

R³ is Dpr, 4PyrAla or (L)- or (D)- Arg;

R⁴ is HomArg, Orn or Lys;


R⁵ is (D)Gln or (L)- or (D)- Trp;

R⁶ is (L)- or (D)- Gln or (p-NO₂)Phe; and

Y² is amide, thioether, thioester or disulfide.

21. A method for treating disorders selected from the group consisting of neoplasms, bacterial, parasite and viral infections, chronic autoimmune disorders and osteoporosis, comprising administering to a mammal in need thereof a pharmaceutical composition comprising a therapeutically effective amount of a backbone cyclized IL-6 antagonist.

22. The method of claim 21 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the general formula 1:



Formula No. 1

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R²⁴⁹ is Trp, (L) or (D)Lys, (L) or (D) Tyr or (D)Phe;

R²⁵⁰ is Arg;

R²⁵¹ is (L) or (D)Leu or Lys;

R²⁵² is (L) or (D)Arg;

R²⁵³ is (D)- or (L)- Phe;

R²⁵⁴ is Ala;

R²⁵⁵ is (D)- or (L)- Leu or is Lys;

R²⁵⁶ is absent or is (L) or (D) Arg;

R²⁵⁷ is (L) or (D) Tyr;

R²⁵⁸ is Ala; and


Y² is amide, thioether, thioester or disulfide.

23. The method of claim 22 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:
Trp-Arg-Lys-(D)Arg-Phe-AlaC3-Leu-Arg-(D)Tyr-AlaN3-NH₂

24. The method of claim 22 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:
(D)Lys-Arg-(D)Leu-(D)Arg-(D)Phe-AlaC3-(D)Leu-Arg-(D)Tyr-AlaN3-NH₂

25. The method of claim 22 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the formula:
(D)Phe-Arg-(D)Leu-(D)Arg-(D)Phe-AlaC3-Leu-(D)Tyr-AlaN3-NH₂

26. The method of claim 21 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the general formula 3:



Formula No. 3

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R¹ is (D)Bip, Gln, Lys, Lys(ZCL) or Dab;

R² is (D)Lys, Gly, Ala or Trp

R³ is Orn, 4PyrAla, (L) or (D)Dab, (D)Arg, Lys or Dpr;

R⁴ is Lys, Lys(ZCL), Arg, Arg(Mtr) or (D)Glu;


R⁵ is Asn, Trp or (D)Ala;

R⁶ is Arg, (p-NO₂)Phe, (L)- or (D)- Trp, Gln, Abu or Glu;

and

Y² is amide, thioether, thioester or disulfide.

27. The method of claim 21 wherein the IL-6 antagonist is a backbone cyclized peptide analog having the general formula 4:



Formula No. 4

wherein m and n are 1 to 5;

X designates a terminal carboxy acid, amide or alcohol group;

R¹ is (D)Phe or Lys;

R² is (D)Cit, Lys or (D)Bip;

R³ is Dpr, 4PyrAla or (L)- or (D)- Arg;

R⁴ is HomArg, Orn or Lys;

R⁵ is (D)Gln or (L)- or (D)- Trp;

R⁶ is (L)- or (D)- Gln or (p-NO₂)Phe; and

Y² is amide, thioether, thioester or disulfide.

28. The method of claim 21 wherein the disorder is selected from the group consisting of rheumatoid arthritis, multiple myeloma and osteoporosis.

Add
B9